arXiv:1409.1610v3 [hep-ph] 1 Jun 2015

Neutrino Oscillations With Recently Measured Sterile-Active Neutrino Mixing Angle

Leonard S. Kisslinger
Department of Physics, Carnegie Mellon University, Pittsburgh, PA 15213

June 2, 2015

Keywords:Neutrino oscillations; Sterile neutrinos; New mixing angle
PACS Indices:11.30.Er,14.60.Lm,13.15.+g

Abstract

This brief report is an extension of a prediction of neutrino oscillation with a sterile neutrino using parameters of the sterile neutrino mass and mixing angle recently extracted from experiment.

1 Introduction

Recently there was an investigation of how the neutrino transition probability $\mathcal{P}(\nu_\mu \to \nu_e)$ is modified by the introduction of a sterile neutrino[1], motivated in part by recent experiments on neutrino oscillations[2], which suggested the existence of at least one sterile neutrino. The 3x3 U-matrix method of Sato and collaborators for three active neutrino oscillations[3, 4] was extended to 4x4 with a sterile neutrino.

In the present brief report we use the formalism of Ref[1] with parameters for the sterile neutrino that have recently been extracted by a global fit to neutrino oscillation data[5].

2 $\mathcal{P}(\nu_\mu \to \nu_e)$ Using a 4x4 U Matrix

Active neutrinos with flavors ν_e, ν_μ, ν_τ and a sterile neutrino ν_s are related to neutrinos with definite mass by

$$\nu_f \;\; = \;\; U \nu_m \; , \tag{1}$$

where U is a 4x4 matrix and ν_f, ν_m are 4x1 column vectors, which is an extension of the 3x3 matrix used in Refs.[3, 4].

As shown in Ref[1], the transition probability $\mathcal{P}(\nu_\mu \to \nu_e)$, assuming $\delta_{CP} = 0$ giving $U^*_{ij} = U_{ij}$, is

$$\begin{aligned}\mathcal{P}(\nu_\mu \to \nu_e) &= U^2_{11}U^2_{21} + U^2_{12}U^2_{22} + U^2_{13}U^2_{23} + \\ &\quad U^2_{14}U^2_{24} + 2U_{11}U_{21}U_{12}U_{22}cos\delta L + \\ &\quad 2(U_{11}U_{21}U_{13}U_{23} + U_{12}U_{22}U_{13}U_{23})cos\Delta L + \\ &\quad 2U_{14}U_{24}(U_{11}U_{21} + U_{12}U_{22} + U_{13}U_{23})cos\gamma L\ , \end{aligned} \tag{2}$$

with neutrino mass differences $\delta m^2_{ij} = m^2_i - m^2_j$, $\delta = \delta m^2_{12}/2E$, $\Delta = \delta m^2_{13}/2E$, $\gamma = \delta m^2_{j4}/2E$ (j=1,2,3), where E is the energy and L the baseline. The neutrino mass differences are $\delta m^2_{12} = 7.6 \times 10^{-5}(eV)^2$, $\delta m^2_{13} = 2.4 \times 10^{-3}(eV)^2$, and from recent MiniBooNE analysis $\delta m^2_{j4} = 0.9(eV)^2$. Note that in Ref[1] there is a typo in the third line of Eq(2), with $U_{13}U_{22}$ rather than the correct $U_{13}U_{23}$.

Using $c_{12} = .83$, $s_{12} = .56$, $s_{23} = c_{23} = .7071$, and $s_{13} = .15$, (with $s_{ij}, c_{ij} = sin\theta_{ij}, cos\theta_{ij}$),

$$\begin{aligned} U_{11} &= .822c_\alpha \\ U_{12} &= .554c_\alpha - .821s^2_\alpha \\ U_{13} &= -.821s^2_\alpha c_\alpha - .554s^2_\alpha + .15c_\alpha \\ U_{14} &= .821s_\alpha c^2_\alpha + .554s_\alpha c_\alpha + .15s_\alpha \\ U_{21} &= -.484c_\alpha \\ U_{22} &= .484s^2_\alpha + .527c_\alpha \\ U_{23} &= .699c_\alpha - (-.484s_\alpha c_\alpha + .527s_\alpha)s_\alpha \\ U_{24} &= -.484s_\alpha c^2_\alpha + .527s_\alpha c_\alpha + .699s_\alpha\ , \end{aligned} \tag{3}$$

with α the sterile-active neutrino mixing angle, $s_\alpha, c_\alpha = sin(\alpha), cos(\alpha)$

A recent analysis of neutrino oscillation data[5] found $sin(\alpha) \simeq 0.16$. Using this, from Eqs(2,3) we obtain the results shown in the figure. The 3x3 results with only active neutrinos are somewhat different than those of Ref[6] as the current value of $s_{13} = .15$ from the Daya Bay experiment[7, 8] was not known at that time.



Figure 1: The ordinate is $\mathcal{P}(\nu_\mu \to \nu_e)$ for MINOS(L=735 km), MiniBooNE(L=500m), T2K(L=295 km), and Double CHOOZ(L=1.03 km) using the 4x4 U matrix with $\delta m^2_{j4} = 0.9(eV)^2$ and $sin(\alpha) \simeq 0.16$. The dashed curves are for $\alpha = 0$ (3x3).

3 Conclusions

As shown in the figure, the neutrino oscillation probability, $\mathcal{P}(\nu_\mu \rightarrow \nu_e)$, is quite different for a model with four neutrinos, $\nu_e, \nu_\mu, \nu_\tau, \nu_s$ than with three active neutrinos. The effect of the sterile neutrino is largest for large values of E/L, as seen in the figure for MiniBooNE, which has a baseline of L=0.5 km. Although Double CHOOZ has a baseline of only 1.03 km, the energy is only about 3 MeV, so the effects of a sterile neutrino are smaller, but are still important for the extraction of neutrino parameters from neutrino oscillation data.

In the present work we have used the sterile-active neutrino mixing angle α ($sin(\alpha) \simeq 0.16$), which is now known to be approximately correct for $\delta m^2_{j4} = 0.9(eV)^2$, rather than the mixing angles used in Ref[1]. Therefore the present results are more reliable for comparison with future neutrino oscillation experiments, inlcuding measurements of CP and T-reversal violations, than the previous publication.

Acknowledgements

This work was carried out while the author was a visitor at Los Alamos National Laboratory, Group P25, and the author thanks Dr. William C Louis III for information on sterile neutrino parameters from MiniBooNE data and several helpful discussions.

References

[1] Leonard S. Kisslinger, arXiv1309.4983, Int. J. Theor.Phys. **53**, 3201 (2014)

[2] A.A. Aguilar-Arevalo *et. al.* (MiniBooNE Collaboration), Phs. Rev. Lett. **110**, 161801 (2013)

[3] Jiro Arafune and Joe Sato, Phys. Rev. **D 55**, 1653 (1997)

[4] Masafumi Koike and Joe Sato, arXiv:hep-ph/9707203 (1997)

[5] Joachim Kopp, Pedro A.N. Machado, Michele Maltoni, Thomas Schwetz, JHEP 1305:050 (2013)

[6] L.S. Kisslinger, E.M. Henley, and M.B. Johnson, Int J. Mod. Phys. E **21**, 1250065 (2012)

[7] F.P. An et.al., Daya Bay Collaboration, arXiv:1210.6327

[8] F.P. An et.al., Chin. Phys. **C37**, 01101 (2013)